

Generation of two-photon EPR and W states

W.B. Cardoso,¹ W.C. Qiang,² A.T. Avelar,¹ and B. Baseia¹

¹*Instituto de Física, Universidade Federal de Goiás, 74.001-970, Goiânia, Goiás, Brazil*

²*Faculty of Science, Xi'an University of Architecture and Technology, 710055, Xi'an, China*

In this paper we present a scheme for generation of two-photon EPR and W states in the cavity QED context. The scheme requires only one three-level Rydberg atom and two or three cavities. The atom is sent to interact with cavities previously prepared in vacuum states, via two-photon process. An appropriate choice of the interaction times allows one to obtain the mentioned states with maximized fidelities. These specific times and the values of success probability and fidelity are discussed.

The entanglement is present in a diversity of applications of quantum mechanics, such as quantum teleportation [1], quantum computation [2], one-way quantum computation [3], quantum communication via teleportation [4, 5], quantum metrology [6], and quantum information escaping from black holes [7]. There are various kinds of entangled states depending on the number of involved parties and of the entanglement 'structure', e.g., for bipartite entanglements there is only one example, the EPR state [8]; for 3-partite there are two kinds of entangled states, the GHZ [9] and W [10] states; for 4-partite, more interesting is the 4-qubit cluster state whose correlations cannot be described in terms of the local realism as experimentally demonstrated in [11]. In fact, this state is not biseparable and has a genuine 4-qubit entanglement.

With respect to the tripartite entangled states, it is known that W states are robust against losses of qubits since they retain bipartite entanglement if we trace out any one qubit, whereas GHZ states are fragile since the remaining two partite states result in separable states. This property turns the W states very attractive for various quantum information tasks, such as a quantum channel for teleportation of entangled pairs [12], probabilistic teleportation [13] and quantum key distribution [14]; however, one should manipulate it with care since it is not appropriate for perfect quantum teleportation and superdense coding in the form presented in Refs.[15, 16]. Several schemes have been proposed to generate the W state in QED cavities [17], trapped ions [18], quantum dots [19], superconducting quantum interference devices [20] and superconducting flux qubits [21].

A class of W states that can be used for perfect teleportation and superdense coding was recently presented in [22], given by

$$|W_\zeta\rangle = \frac{1}{\sqrt{2+2\zeta}}(|001\rangle + \sqrt{\zeta}e^{i\gamma}|010\rangle + \sqrt{\zeta+1}e^{i\delta}|100\rangle), \quad (1)$$

where ζ is a real number and γ, δ are phases. Additionally, in [23] Li and Qiu generalized the states of the W-class for multi-qubit systems and multi-particle systems with higher dimension.

In view of applications of the entangled states discussed above, in this work we present a scheme for generation of two-photon EPR and W states in QED cavities. To this end, a Rydberg three-level atom is sent to interact with the cavities previously prepared in the vacuum state via a two-photon process. It is worth to mention that the two-photon process has been demonstrated in [24] for microwave cavity QED and

offers some advantages in relation to one-photon process, as the reduction of interaction times due to the increasing of the atom-field coupling strength and lower decoherence induced by stray fields [25]. In addition, two-photon process can be easily obtained with Rydberg atoms with principal quantum number $n > 89$, which can be state-sensitively detected using tunneling field ionization with quantum efficiencies above 80% and an ionization efficiency above 98% [26]. In the following, we discuss the theoretical model and give details of our scheme.

Consider a three-level atom that interacts with a single cavity-field mode via two-photon Jaynes-Cummings model described in the interaction picture by the Hamiltonian [27]

$$H_I = \hbar g_1 (a|e\rangle\langle f|e^{-i\delta t} + a^\dagger|f\rangle\langle e|e^{i\delta t}) + \hbar g_2 (a|f\rangle\langle g|e^{i\delta t} + a^\dagger|g\rangle\langle f|e^{-i\delta t}), \quad (2)$$

where g_1 and g_2 stand for the one-photon coupling constant with respect to the transitions $|e\rangle \leftrightarrow |f\rangle$ and $|f\rangle \leftrightarrow |g\rangle$, respectively. The detuning δ is given by

$$\delta = \Omega - (\omega_e - \omega_f) = (\omega_f - \omega_g) - \Omega, \quad (3)$$

where Ω is the cavity-field frequency and ω_e, ω_f , and ω_g are the frequencies associated with the atomic levels $|e\rangle, |f\rangle$, and $|g\rangle$, respectively. Fig. 1 shows a schematic representation of the atomic levels.

The state describing the combined atom-field system reads

$$|\psi(t)\rangle = \sum_n [C_{e,n}(t)|e, n\rangle + C_{f,n}(t)|f, n\rangle + C_{g,n}(t)|g, n\rangle], \quad (4)$$

where $|k, n\rangle$, with $k = e, f, g$, indicating the atom in the state $|k\rangle$ and the field in the Fock state $|n\rangle$. The coefficients $C_{k,n}(t)$ stand for the corresponding probability amplitudes.




Figure 1: Schematic diagram of the three-level atom interacting with a single-mode of a cavity field.

Inserting the Eqs. (2) and (4) into the time dependent Schrödinger equation one obtains the coupled first-order differential equations for the probability amplitudes

$$\begin{aligned}\frac{dC_{e,n}(t)}{dt} &= -ig_1C_{f,n+1}(t)\sqrt{n+1}e^{-i\delta t}, \\ \frac{dC_{f,n+1}(t)}{dt} &= -ig_1C_{e,n}(t)\sqrt{n+1}e^{i\delta t} \\ &\quad -ig_2C_{g,n+2}(t)\sqrt{n+2}e^{i\delta t}, \\ \frac{dC_{g,n+2}(t)}{dt} &= -ig_2C_{f,n+1}(t)\sqrt{n+2}e^{-i\delta t}.\end{aligned}\quad (5)$$

As usually, we consider that the entire atom-field system is

decoupled at the initial time $t = 0$,

$$\begin{aligned}C_{e,n}(0) &= C_e C_n(0), \\ C_{f,n+1}(0) &= C_f C_{n+1}(0), \\ C_{g,n+2}(0) &= C_g C_{n+2}(0),\end{aligned}\quad (6)$$

where the $C_n(0)$ stand for the amplitudes of the arbitrary initial field state and the C_a are atomic amplitudes of the (normalized) initial atomic state

$$|\chi\rangle = C_e|e\rangle + C_f|f\rangle + C_g|g\rangle. \quad (7)$$

Solving these coupled differential equations with the initial conditions in (6) we get the time dependent coefficients as

$$\begin{aligned}C_{e,n}(t) &= \left[\frac{g_1^2(n+1)}{\Lambda_n \alpha_n^2} \gamma_n(t) + 1 \right] C_e C_n - i \frac{g_1 \sqrt{n+1}}{\Lambda_n} \sin(\Lambda_n t) e^{-i\frac{\delta t}{2}} C_f C_{n+1} \\ &\quad + \left[\frac{g_1 g_2 \sqrt{(n+1)(n+2)}}{\Lambda_n \alpha_n^2} \gamma_n(t) \right] C_g C_{n+2},\end{aligned}\quad (8)$$

$$\begin{aligned}C_{f,n+1}(t) &= -i \frac{g_1 \sqrt{n+1}}{\Lambda_n} \sin(\Lambda_n t) e^{i\frac{\delta t}{2}} C_e C_n + \left(\cos(\Lambda_n t) - \frac{i\delta}{2\Lambda_n} \sin(\Lambda_n t) \right) e^{i\frac{\delta t}{2}} C_f C_{n+1} \\ &\quad - i \frac{g_2 \sqrt{n+2}}{\Lambda_n} \sin(\Lambda_n t) e^{i\frac{\delta t}{2}} C_g C_{n+2},\end{aligned}\quad (9)$$

$$\begin{aligned}C_{g,n+2}(t) &= \frac{g_1 g_2 \sqrt{(n+1)(n+2)}}{\Lambda_n \alpha_n^2} \gamma_n(t) C_e C_n - i \frac{g_2 \sqrt{n+2}}{\Lambda_n} \sin(\Lambda_n t) e^{-i\frac{\delta t}{2}} C_f C_{n+1} \\ &\quad + \left[\frac{g_2^2(n+2)}{\Lambda_n \alpha_n^2} \gamma_n(t) + 1 \right] C_g C_{n+2},\end{aligned}\quad (10)$$

where

$$\gamma_n(t) = \left[\Lambda_n \cos(\Lambda_n t) + i \frac{\delta}{2} \sin(\Lambda_n t) - \Lambda_n e^{i\frac{\delta t}{2}} \right] e^{-i\frac{\delta t}{2}}, \quad (11)$$

$$\Lambda_n = \sqrt{\frac{\delta^2}{4} + \alpha_n^2}, \quad (12)$$

$$\alpha_n = \sqrt{g_1^2(n+1) + g_2^2(n+2)}, \quad (13)$$

Λ_n being the Rabi frequency. The substitutions $n \rightarrow n-1$ in Eq. (9) and $n \rightarrow n-2$ in Eq. (10) allow one to obtain the $C_{f,n}(t)$ and $C_{g,n}(t)$, respectively.

Two-photon EPR state - Now, we describe the scheme for generation of the maximally entangled two-photon EPR state, written in the form

$$|\psi\rangle_{12} = \frac{1}{\sqrt{2}}(|02\rangle_{12} + |20\rangle_{12}), \quad (14)$$

where the subscripts 1 and 2 concern the cavities 1 and 2, respectively. Assume the two cavities previously prepared in the vacuum state and the atom in its excited state ($|e\rangle$).

Firstly, the atom crosses the cavity 1, interacting with the field mode for a time t_1 . Thus, the initial state describing the atom plus the two cavities evolves to

$$\begin{aligned}|\phi\rangle_{a12} &= C_{e0}^{(e0)}(t_1)|e00\rangle_{a12} + C_{f1}^{(e0)}(t_1)|f10\rangle_{a12} \\ &\quad + C_{g2}^{(e0)}(t_1)|g20\rangle_{a12},\end{aligned}\quad (15)$$

where $C_{ij}^{(k,l)}$ are the coefficients in Eqs. (8)-(10) considering the input atomic state in k and the field state in l .

Next, the atom enters the cavity 2 and interacts for a time t_2 . So, the state of the whole system is written as

$$\begin{aligned}|\phi'\rangle_{a12} &= C_{e0}^{(e0)}(t_1)[C_{e0}^{(e0)}(t_2)|e00\rangle_{a12} + C_{f1}^{(e0)}(t_2)|f01\rangle_{a12} \\ &\quad + C_{g2}^{(e0)}(t_2)|g02\rangle_{a12}] + C_{f1}^{(e0)}(t_1)[C_{f0}^{(f0)}(t_2)|f10\rangle_{a12} \\ &\quad + C_{g1}^{(f0)}(t_2)|g11\rangle_{a12}] + C_{g2}^{(e0)}(t_1)|g20\rangle_{a12}.\end{aligned}\quad (16)$$




Figure 2: (Color online) Plots of (a) success probability of detection of the atomic state in $|g\rangle$ and (b) the respective fidelity for the two-photon EPR state, for parameters $g_1 = g_2 = g = 17.5\text{MHz}$, $\delta = 30g$.




Figure 3: (Color online) Comparison of the fidelity and success probability versus the time t_2 . We use the values $t_1 = 2\mu\text{s}$ for the fidelity in dashed line (green) and for success probability in dotted line (red) and $t_1 = 5\mu\text{s}$ for the fidelity in solid line (blue) and for the success probability in dash-dot line (black), with same conventions of Fig 2.

To get the two-photon EPR state we may choose to realize the atomic detection or not. In the affirmative case, the detection of the atomic state $|g\rangle$ leads to the success probability

$$P_g = |C_{e0}^{(e0)}(t_1)|^2 |C_{g2}^{(e0)}(t_2)|^2 + |C_{f1}^{(e0)}(t_1)|^2 |C_{g1}^{(f0)}(t_2)|^2 + |C_{g2}^{(e0)}(t_1)|^2, \quad (17)$$

and the respective fidelity

$$F = \frac{1}{2P_g} |C_{e0}^{(e0)}(t_1)C_{g2}^{(e0)}(t_2) + C_{g2}^{(e0)}(t_1)|^2. \quad (18)$$

The success probability and fidelity of the EPR state with atomic detection is displayed in Fig. 2a and 2b, respectively. The values of experimental parameters $g_1 = g_2 = g = 17.5\text{MHz}$ and $\delta = 30g$ were used. In this case, Fig. 3 presents the values of the success probability and fidelity for two different times. Note that an appropriate choice of the interaction time allows one to obtain a greater fidelity.

On the other hand, if we choose to realize no atomic detection, the maximum value of the fidelity decreases to a value close to 0.8, with the form

$$F = \frac{1}{2} |C_{e0}^{(e0)}(t_1)C_{g2}^{(e0)}(t_2) + C_{g2}^{(e0)}(t_1)|^2, \quad (19)$$

as used in the Fig. 4.

Two-photon W state - The W state to be constructed is written as

$$|\psi\rangle_{123} = \frac{1}{2}(|002\rangle_{123} + |020\rangle_{123} + \sqrt{2}|200\rangle_{123}), \quad (20)$$




Figure 4: (Color online) Fidelity of the two-photon EPR state without atomic detection, using the same conventions of Fig 2.

where the subscripts concern the cavities 1, 2, and 3, respectively. Also, we can generate the two-photon W state using a similar procedure to the two-photon EPR state generation. However, the W state is realized in three cavities and a distinct procedure is required.

Firstly, we consider these three cavities in the vacuum state. An atom in the excited state $|e\rangle$ is sent to interact with the first cavity by a time t_1 , leading the whole system in the state

$$|\phi\rangle_{a123} = C_{e0}^{(e0)}(t_1)|e000\rangle_{a123} + C_{f1}^{(e0)}(t_1)|f100\rangle_{a123} + C_{g2}^{(e0)}(t_1)|g200\rangle_{a123}. \quad (21)$$

Next, the atom crosses the cavities 2 and 3 by a time t_2 and t_3 , respectively. The above state goes to the form

$$\begin{aligned} |\phi''\rangle_{a123} = & C_{e0}^{(e0)}(t_1)\{C_{e0}^{(e0)}(t_2)[C_{e0}^{(e0)}(t_3)|e000\rangle_{a123} \\ & + C_{f1}^{(e0)}(t_3)|f001\rangle_{a123} + C_{g2}^{(e0)}(t_3)|g002\rangle_{a123}\} \\ & + C_{f1}^{(e0)}(t_2)[C_{f0}^{(f0)}(t_3)|f010\rangle_{a123} \\ & + C_{g1}^{(f0)}(t_3)|g011\rangle_{a123} + C_{g2}^{(e0)}(t_2)|g020\rangle_{a123}\} \\ & + C_{f1}^{(e0)}(t_1)\{C_{f0}^{(f0)}(t_2)[C_{f0}^{(f0)}(t_3)|f100\rangle_{a123} \\ & + C_{g1}^{(f0)}(t_3)|g101\rangle_{a123} + C_{g1}^{(f0)}(t_2)|g110\rangle_{a123}\} \\ & + C_{g2}^{(e0)}(t_1)|g200\rangle_{a123}. \end{aligned} \quad (22)$$

If we detect the atom in its ground state ($|g\rangle$) the state of the system collapses in the form

$$\begin{aligned} |\phi'''\rangle_{123} = & N\{C_{e0}^{(e0)}(t_1)[C_{e0}^{(e0)}(t_2)C_{g2}^{(e0)}(t_3)|002\rangle_{123} \\ & + C_{f1}^{(e0)}(t_2)C_{g1}^{(f0)}(t_3)|011\rangle_{123} + C_{g2}^{(e0)}(t_2)|020\rangle_{123}\} \\ & + C_{f1}^{(e0)}(t_1)[C_{f0}^{(f0)}(t_2)C_{g1}^{(f0)}(t_3)|101\rangle_{123} \\ & + C_{g1}^{(f0)}(t_2)|110\rangle_{123} + C_{g2}^{(e0)}(t_1)|200\rangle_{123}, \end{aligned} \quad (23)$$

with success probability given by

$$\begin{aligned} P_g = & |C_{e0}^{(e0)}(t_1)|^2 |C_{e0}^{(e0)}(t_2)|^2 |C_{g2}^{(e0)}(t_3)|^2 \\ & + |C_{e0}^{(e0)}(t_1)|^2 |C_{f1}^{(e0)}(t_2)|^2 |C_{g1}^{(f0)}(t_3)|^2 \\ & + |C_{e0}^{(e0)}(t_1)|^2 |C_{g2}^{(e0)}(t_2)|^2 \\ & + |C_{f1}^{(e0)}(t_1)|^2 |C_{f0}^{(f0)}(t_2)|^2 |C_{g1}^{(f0)}(t_3)|^2 \\ & + |C_{f1}^{(e0)}(t_1)|^2 |C_{g1}^{(f0)}(t_2)|^2 + |C_{g2}^{(e0)}(t_1)|^2. \end{aligned} \quad (24)$$




Figure 5: (Color online) Plots of fidelity and success probability versus the interaction time t_3 for the two-photon W state generation scheme. In (a) we consider the atomic detection in the state $|g\rangle$, with the values $t_1 = t_2 = 4\mu\text{s}$ for the fidelity in dotted line (red) and for success probability in dashed line (green) and the values $t_1 = t_2 = 7\mu\text{s}$ for the fidelity in dash-dot line (black) and for success probability in solid line (blue). In (b) we consider the scheme without atomic detection. For this case we show three values of the fidelity versus t_3 . In solid line (blue) we use $t_1 = t_2 = 16\mu\text{s}$; in dotted line (red) we use $t_1 = t_2 = 2\mu\text{s}$; in dashed line (green) we use $t_1 = t_2 = 13\mu\text{s}$, using the same conventions of Fig 2.

We calculate the fidelity $F = |\langle \psi | \phi \rangle|_{123}^2$, written as

$$F = \frac{1}{4Pg} |C_{e0}^{(e0)}(t_1)C_{e0}^{(e0)}(t_2)C_{g2}^{(e0)}(t_3) + C_{e0}^{(e0)}(t_1)C_{g2}^{(e0)}(t_2) + \sqrt{2}C_{g2}^{(e0)}(t_1)|^2. \quad (25)$$

With an appropriate choice of t_i ($i = 1, 2, 3$) one gets the maximized fidelity. For example, in the case of atomic detections we choose the interaction times $t_1 = t_2 = t_3 = 32\mu\text{s}$ to obtain approximately success probability 30% and fidelity 95%.

In conclusion, we presented a scheme for the generation of two-photon EPR and W states in the cavity QED context. Concerning with the experimental feasibility, the devices used here were based on Ref. [25] concerned with Rydberg atoms with quantum number $n \sim 90$. The coupling constant of these atoms with the cavity is taken close to $g = 17.5\text{MHz}$. In our simulations we have used this value and a detuning of $30g$. As example, by choosing the interaction times as $t_1 = t_2 = 3\mu\text{s}$ for the two-photon EPR state the success probability and fidelity result approximately 40% and 97%, respectively; if we choose $t_1 = t_2 = t_3 = 32\mu\text{s}$ for the two-photon W state, the success probability and fidelity result 30% and 95%, respectively. On the other hand, the time spent for the entire procedure is approximately 10^{-4}s for both, EPR and W case. These times are much smaller than the decoherence time of the cavity (0.1s) [28], which shows the experimental feasibility of the scheme.

We thank the CAPES, CNPq, and FUNAPE/GO, Brazilian agencies, for the partial supports. This work is also partially supported by Natural Science Basic Research Plan in the Shaanxi Province of China (program no: SJ08A13), the Natural Science Foundation of the Education Bureau of Shaanxi Province, China under Grant O9jk534.

-
- [1] C.H. Bennett *et al.*, Phys. Rev. Lett. **70**, 1895 (1993).
 - [2] D. Gottesman and I.L. Chuang, Nature **402**, 390 (1999); E. Knill, R. Laflamme, and G.J. Milburn, Nature **409**, 46 (2001).
 - [3] R. Raussendorf and H.J. Briegel, Phys. Rev. Lett. **86**, 5188 (2001).
 - [4] J.I. Cirac *et al.*, Phys. Rev. Lett. **78**, 3221 (1997), and references therein.
 - [5] G. Brassard, S.L. Braunstein, and R. Cleve, Physica D **120**, 43 (1998).
 - [6] A. Gilchrist *et al.*, J. Opt. B: Quantum Semiclass. Opt. **6**, S828 (2004).
 - [7] S. Lloyd, Phys. Rev. Lett. **96**, 061302 (2006).
 - [8] H.K. Lo and S. Popescu, Phys. Rev. A **63**, 022301 (2001).
 - [9] W. Dur, G. Vidal, and J.I. Cirac, Phys. Rev. A **62**, 062314 (2000).
 - [10] A. Acin *et al.*, Phys. Rev. Lett. **87**, 040401 (2001).
 - [11] P. Walther *et al.*, Phys. Rev. Lett. **95**, 020403 (2005).
 - [12] V.N. Gorbachev *et al.*, Phys. Lett. A **314**, 267 (2003).
 - [13] J. Joo *et al.*, New J. Phys. **5**, 136 (2003).
 - [14] J. Joo *et al.*, arXiv:quant-ph/0204003 (2002).
 - [15] A. Karlsson and M. Bourennane, Phys. Rev. A **58**, 4394 (1998).
 - [16] A.K. Pati, Phys. Rev. A **61**, 022308 (2000).
 - [17] G.P. Guo *et al.*, Phys. Rev. A **65**, 042102 (2002); G.C. Guo and Y.S. Zhang, Phys. Rev. A **65**, 054302 (2002); Z.J. Deng, M. Feng, and K.L. Gao, Phys. Rev. A **73**, 014302 (2006); E. M. Bécerra-Castro *et al.*, J. Phys. B: At. Mol. Opt. Phys. **41**, 215505 (2008).
 - [18] G.X. Li, Phys. Rev. A **74**, 055801 (2006); W.X. Yang, J.H. Li, and A.S. Zheng, Commun. Theor. Phys. **47**, 421 (2007).
 - [19] X. Wang, M. Feng, and B.C. Sanders, Phys. Rev. A **67**, 022302 (2003).
 - [20] A.S. Zheng *et al.*, Chin. Phys. Lett. **25**, 1195 (2007).
 - [21] M.D. Kim and S.Y. Cho, Phys. Rev. B **77**, 100508(R) (2008).
 - [22] P. Agrawal and A. Pati, Phys. Rev. A **74**, 062320 (2006).
 - [23] L. Li and D. Qiu, J. Phys. A: Math. Theor. **40**, 10871 (2007).
 - [24] M. Brune *et al.*, Phys. Rev. Lett. **59**, 1899 (1987).
 - [25] P. J. Blythe and B. T. H. Varcoe, New J. Phys. **8**, 213 (2006).
 - [26] M. Tada *et al.*, Phys. Lett. A **303**, 285 (2002); Y. Kishimoto *et al.*, Phys. Lett. A **303**, 279 (2002).
 - [27] A.H. Toor and M.S. Zubairy, Phys. Rev. A **45**, 4951 (1992); A.D. dSouza *et al.*, Physica A **388**, 1331 (2009).
 - [28] S. Gleyzes *et al.*, Nature **446**, 297 (2007); J. M. Raimond, M. Brune, and S. Haroche, Rev. Mod. Phys. **73**, 565 (2001).